Kafkas Universitesi Veterinar Fakultesi Dergisi
Journal Home-Page: http://vetdergikafkas.org
E-ISSN: 1309-2251

Kafkas Univ Vet Fak Derg
29 (3): 247-254, 2023
DOI: 10.9775/kvfd.2023.29134

RESEARCH ARTICLE

The Effect of *Bacillus subtilis* and Fructooligosaccharide as Antibiotic Substituent on Goose Performance Parameters, Serum Biochemical Indicators and Intestinal Morphology

Guang-quan LI [1 (‡)] Xian-ze WANG [1 (‡)] Yi LIU [1] Yun-zhou YANG [1] Cui WANG [1] Shao-ming GONG [1] Da-qian HE [1 (*)] Hui-ying WANG [1 (*)]

(‡) These authors contributed equally to this study

[1] Institute of Animal Husbandry and Veterinary Science, Shanghai Academy of Agricultural Science, Shanghai, 201106, P. R. CHINA

ORCIDs: G.Q.L. 0000-0003-1050-634X; X.Z.W. 0000-0002-7507-3437; Y.L. 0000-0002-7237-6010; Y.Z.Y. 0000-0003-1036-8047; C.W. 0000-0003-2723-5929; S.M.G. 0000-0002-9354-1804; D.Q.H. 0000-0002-8714-4781; H.Y.W. 0000-0002-8312-7028

Article ID: KVFD-2023-29134 Received: 05.02.2023 Accepted: 02.05.2023 Published Online: 10.05.2023

Abstract: As antibiotics are now prohibited or heavily regulated in farming in most countries, nutritionists and veterinarians are increasingly turning to probiotic supplements, prebiotics, and synbiotics to promote animal health and improve production performance. The study aimed to evaluate the effects of *Bacillus subtilis*, fructooligosaccharides (FOS), and their combined use on geese, compared to antibiotics, and evaluate their suitability as an alternative to antibiotics. 240 14-day-old geese were randomly divided into five groups and were fed a basal diet or supplemented with 100 mg/kg chlortetracycline, 400 mg/kg FOS, 2x10^{5}cfu g/kg *B. subtilis*, or a combination of 200 mg/kg FOS and 1x10^{5}cfu g/kg *B. subtilis* for 56 days. The study measured the body weight, feed consumption, serum parameters, and intestinal morphology of the geese. The results showed that the combination of *B. subtilis* and FOS had a positive impact on the body weight, feed consumption, and serum parameters of the geese, while improving their intestinal morphology. The results suggest that the combination of *B. subtilis* and FOS may provide multiple benefits to animal health and performance and could be used as a suitable alternative to antibiotics.

Keywords: *Bacillus subtilis, Fructooligosaccharide, Goose, Intestinal morphology, Performance parameters, Serum*

Antibiyotik İkamesi Olarak *Bacillus subtilis* ve Fruktooligosakkaritin Kazlarda Performans Parametreleri, Serum Biyokimyasal Göstergeleri ve Bağırsak Morfolojisi Üzerine Etkisi

Öz: Antibiyotikler artık çoğu ülkede çiftçilikte yasaklandığından veya ağır düzenlemelere tabi tutulduğundan, beslenme uzmanları ve veterinerler hayvan sağlığını desteklemek ve üretim performansını artırmak için probiyotik takviyelere, prebiyotiklere ve sinbiyotiklere giderek daha fazla yönelmektedir. Bu çalışmanın amacı, antibiyotiklere kıyasla *Bacillus subtilis*, fruktooligosakkaritler (FOS) ve bunların birlikte kullanımının kazlar üzerindeki etkilerini ve antibiyotiklere alternatif olarak uygunluğunu değerlendirmektir. İkiyüz kırk adet 14 günlük kaz rastgele beş gruba ayrılmış ve 56 gün boyunca bazal diyet veya 100 mg/kg klortetrasiklin, 400 mg/kg FOS, 2x10^{5}cfu g/kg *B. subtilis* ve 200 mg/kg FOS ve 1x10^{5}cfu g/kg *B. subtilis* kombinasyonu ile beslenmiştir. Çalışmada kazların vücut ağırlığı, yem tüketimi, serum parametreleri ve bağırsak morfolojisi ölçülmüştür. Sonuçlar, *B. subtilis* ve FOS kombinasyonunun kazların vücut ağırlığı, yem tüketimi ve serum parametreleri üzerinde olumlu bir etkiye sahip olduğunu ve bağırsak morfolojilerini iyileştirdiğini göstermiştir. Sonuçlar, *B. subtilis* ve FOS kombinasyonunun hayvan sağlığı ve performansına birçok fayda sağlayabileceğini ve antibiyotiklere uygun bir alternatif olarak kullanılabileceğini göstermektedir.

Anahtar sözcükler: *Bacillus subtilis, Bağırsak morfolojisi, Fruktooligosakkarit, Kaz, Performans parametreleri*

How to cite this article?

Li GQ, Wang XZ, Liu Y, Yang YZ, Wang C, Gong SM, He DQ, Wang HY: The effect of *Bacillus subtilis* and fructooligosaccharide as antibiotic substituent on goose performance parameters, serum biochemical indicators and intestinal morphology. *Kafkas Univ Vet Fak Derg,* 29 (3): 247-254, 2023.
DOI: 10.9775/kvfd.2023.29134

(*) Corresponding author: Da-qian HE & Hui-ying WANG

Phone: +86-18918162216 (D.Q. He); +86-18964198956 (H.Y. Wang)
E-mail: daqianhe@aliyun.com (D.Q. He); yjshywang@sina.com (H.Y. Wang)

This article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

Introduction

Antibiotics have a long history of being added to animal feed as growth promoters [1], It is widely used because of its ability to treat diseases infected by microorganisms such as bacteria, chlamydia, and mycoplasma [2]. However, with the continuous understanding of antibiotics, many bacteria have developed resistance to antibiotics [3], and the abuse of antibiotics will remain in animals or products [4,5], which will damage human health [6].

Probiotics widely exist in nature, animals and plants, can secrete a variety of metabolites, have a wide range of beneficial effects on the host [7]. As a typical probiotic, *B. subtilis* can secrete protease and cellulase to promote the digestion and utilization of nutrients [8]. At the same time, *B. subtilis* can produce spores during the dormant period [9], and has tolerance to high temperature and pressure during feed processing and the acidic environment of the animal gastrointestinal tract [10]. FOS have been widely used in animal husbandry as a prebiotic supplement in animal feed [11,12]. FOS are short chains of fructose molecules that act as a substrate for beneficial bacteria in the gut, promoting the growth of probiotics and maintaining gut health [13]. The application of FOS in animal husbandry has been found to be effective in improving gut microbiota balance, increasing feed efficiency, promoting growth, and reducing the incidence of digestive disorders. FOS have been particularly useful in poultry and swine production, where they have been shown to enhance gut health and improve feed utilization [14,15]. Overall, FOS have been a valuable addition to animal nutrition, providing multiple benefits to animal health and performance.

The use of probiotics or prebiotics alone can have limited benefits when compared to combining them [16,17]. There are few studies on *B. subtilis* and FOS in geese [18]. This experiment aims to study the effects of *B. subtilis*, FOS and their combined use on geese, compare its effect with that of antibiotics, and evaluate whether it can be used as a suitable alternative to antibiotics.

Material and Methods

Animal Ethics

Animal experimentation was approved by the Laboratory Animal Ethics Committee of the Shanghai Academy of Agricultural Sciences (SAASPZ0522046).

Material

The *B. subtilis* used in the experiment (the number of viable bacteria ≥1.0x10^{8}cfu/g) was purchased from Shanghai Shenya Animal Health Products Fuyang Co., Ltd, Anhui, China. FOS (content ≥97.15%) was purchased from Dongguan Zhenshang Industrial Co., Ltd, Guangdong, China. Zhedong white geese were purchased from Xiangshan County Zhejiang White Goose Research Institute, Zhejiang, China.

Experimental Design

Total 240 14-day-old Zhedong white geese were randomly divided into five groups, six replicates in each group, each replicate has eight geese (half male and half female). The five groups were basal diet group, 100mg/kg chlortetracycline (CTC) group, 400 mg/kg FOS group, 2x10^{5}cfu g/kg *B. subtilis* group and 200mg/kg FOS+ 1x10^{5}cfu g/kg *B. subtilis* group, the experiment lasts for 56 d [19]. The basal diet refers to the nutritional level recommended by NRC-1994 (*Table 1*) [20]. During the experiment period, Geese in each replicated was housed in an 80 cm x 80 cm cage with access to natural light during the day and low-level artificial light at night. They had free access to food and water, and routine feeding management and immunization procedures were carried out.

Performance Parameters

Body weight (BW) was recorded before the start and end of the experiment (8 h fasting, 2 h water deprivation), then weigh the amount of feed consumed for each repetition, and the average daily gain (ADG), average daily feed intake (ADFI) and feed/gain (F/G) were calculated at last.

Table 1. *Feed ingredients and analyzed chemical composition of geese diets (air-dry basis %)*

Ingredients	Content %	
	1-28 d	28-70 d
Corn	60.30	58.80
Soybean meal (43% CP)	32.60	25.60
Fish meal (60.3% CP)	2.00	10.10
Soybean oil	2.00	1.50
Lys + Met	0.10	0.00
Limestone	0.00	1.00
Premix [a]	3.00	3.00
Total	100	100
Nutritional Level		
ME/(MJ/kg)	12.13	12.55
Crude protein	20.23	16.00
Crude fiber	3.07	7.00
Ca	0.55	0.68
P	0.45	0.43

[a] *Per kilogram of diets including: Vit-A: 1500 IU, Vit-B_1: 2.3 mg, Vit-B_2: 5.0 mg, Vit-B_6: 5 mg, Vit-B_{12}: 2 mg, Vit-D_3: 200 IU, Vit-E: 12.5 IU, Vit-K: 1.5 mg, Trace elements: 50 g, Garlicin: 30 g, Lysine: 100 g, Methionine: 50 g, Salt: 100 g, Stone powder: 100 g, Myco-Ad: 100 g, Zeolite powder: 420 g;* [b] *Nutrient levels were all calculated values*

Table 2. Effects of B. subtilis and FOS on growth performance in geese

Items	Treatments				
	Basal Diet	100 mg/kg CTC	400 mg/kg FOS	2×10^5cfu g/kg *B. subtilis*	200 mg/kg FOS+1×10^5cfu g/kg *B. subtilis*
Initial BW, g	677.08±14.44	656.47±13.76	659.57±11.23	662.78±13.54	655.93±13.15
Final BW, g	3904.92±103.72^b	4111.11±123.98^a	3919.33±117.51^b	4055.82±85.16^a	4076.97±107.88^a
ADFI, g/d	304.41±1.31	304.96±1.05	301.87±1.06	309.66±1.22	310.03±1.62
ADG, g/d	57.64±8.59^b	61.69±10.26^a	58.21±7.93^{ab}	60.59±8.65^{ab}	61.90±7.86^a
F/G	5.21±0.56^a	4.92±0.86^b	5.13±0.53^{ab}	5.07±0.57^{ab}	5.03±0.93^{ab}

*The data in the table are compared in the same row, and different lowercase letters indicate that the difference has reached a significant level (P<0.05); **CTC:** chlortetracycline, **FOS**: Fructo oligosaccharide*

Serum Biochemical Indicators

At the end of the experiment, one male and female geese with close to average body weight were selected for each repetition, and blood was collected from the wing vein to prepare serum and stored at -20°C for later use. All serum were sent to Shanghai Pinyi Biological Co., Ltd. for testing total protein (TP), albumin (ALB), globulin (GLOB), alanine aminotransferase (ALT), aspartate aminotransferase (AST), glucose (GLU), blood urea nitrogen (BUN), total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C) and low-density lipoprotein cholesterol (LDL-C).

Intestinal Morphology

The geese were slaughtered through exsanguination of the jugular vein.In reference to the methods used by Xue et al.[21], the duodenum, jejunum, and ileum were collected and analyzed. In summary, the contents of their intestines were carefully emptied, rinsed with saline solution, and dried with filter paper. The first quarter of the duodenum, jejunum, and anterior ileum were then cut into 2 cm sections for paraffin embedding. From each sample, five representative intestinal villus crypts were selected, and measurements were taken of the villus height (VH) and crypt depth (CD). The ratio of VH to CD (VH/CD) was calculated.

Statistical Analysis

The data were preliminarily processed using Excel 2019, followed by a one-way ANOVA using SPSS 26.0 software. Multiple comparisons were conducted using Duncan's post hoc test, and the results were presented as mean ± standard deviation. Statistical significance was defined as P<0.05, indicating significant differences.

Results

Performance Parameters

It can be seen from *Table 2* that adding 100 mg/kg CTC to the goose diet was the best for improving the growth performance, compared to the basic diet group, it




Fig 1. The effect of different treatments on goose performance parameters. **CTC:** chlortetracycline, **FOS:** Fructo oligosaccharide, **BW:** body weight, **ADG:** average daily gain, **F/G:** feed/gain




Fig 2. The effect of different treatments on goose serum biochemical indicators. **CTC:** chlortetracycline, **FOS:** Fructo oligosaccharide, **ALT:** alanine aminotransferase, **BUN:** blood urea nitrogen, **HDL-C:** high-density lipoprotein cholesterol




Fig 3. The effect of different treatments on goose intestinal morphology. **CTC:** chlortetracycline, **FOS:** Fructo oligosaccharide

significantly increased the final BW, ADG and F/G ($P<0.05$). The ADG and Final BW of the 200mg/kg FOS+1x10^{5}cfu g/kg *B. subtilis* group improved significantly compared to the basic diet group ($P<0.05$). Adding 2x10^{5}cfu g/kg *B. subtilis* can also improve the Final BW ($P<0.05$), but there is no significant difference in other parameters. There was no significant difference between 400 mg/kg FOS and basal diet group. Significant differences in performance parameters between groups are shown in *Fig. 1*.

Table 3. *Effects of B. subtilis and FOS on serum biochemical indicators in geese*

Items	Treatments				
	Basal Diet	100 mg/kg CTC	400 mg/kg FOS	2×10^{5}cfu g/kg *B. subtilis*	200 mg/kg FOS+1×10^{5}cfu g/kg *B. subtilis*
TP, g/L	31.43±6.48	39.89±3.42	32.90±8.77	37.06±8.83	39.02±7.00
ALB, g/L	12.06±1.97	15.19±1.13	13.74±2.86	14.56±2.67	14.68±2.34
GLOB, g/L	19.36±4.61	24.70±2.50	19.16±5.96	23.50±6.24	24.34±4.67
ALT, U/L	7.50±2.33^b	13.50±3.89^a	7.13±2.53^b	7.75±2.96^b	6.00±2.35^b
AST, U/L	23.25±10.28	21.63±4.47	19.88±7.36	21.13±7.16	20.2±5.12
BUN, mmol/L	0.35±0.15^a	0.14±0.11^b	0.28±0.12^{ab}	0.16±0.09^b	0.18±0.13^b
GLU, mmol/L	6.10±1.45	9.46±1.25	5.50±1.70	6.70±1.26	5.04±2.25
TC, mmol/L	2.74±0.63	3.39±0.69	2.89±1.14	3.75±1.22	2.83±1.08
HDL-C, mmol/L	1.67±0.28^b	2.32±0.44^a	1.70±0.70^b	2.18±0.53^a	2.08±0.60^a
LDL-C, mmol/L	0.98±0.26	0.78±0.17	1.00±0.25	0.76±0.29	0.88±0.50

The data in the table are compared in the same row, and different lowercase letters indicate that the difference has reached a significant level (P<0.05); ***CTC:*** *chlortetracycline,* ***FOS:*** *Fructo oligosac charide,* ***TP:*** *total protein,* ***ALB:*** *albumin,* ***GLOB:*** *globulin,* ***ALT:*** *alanine aminotransferase,* ***AST:*** *aspartate aminotransferase,* ***GLU:*** *glucose,* ***BUN:*** *blood urea nitrogen,* ***TC:*** *total cholesterol,* ***HDL-C:*** *high-density lipoprotein cholesterol,* ***LDL-C:*** *low-density lipoprotein cholesterol*

Table 4. *Effects of B. subtilis and FOS on intestinal morphology in geese*

Intestine	Items	Treatments				
		Basal Diet	100 mg/kg CTC	400 mg/kg FOS	2×10^{5}cfu g/kg *B. subtilis*	200 mg/kg FOS+1×10^{5}cfu g/kg *B. subtilis*
Duodenum	VH, um	504.54±134.18	511.64±66.11	516.74±82.99	504.02±37.84	501.24±66.70
	CD, um	124.03±16.89^a	107.27±37.72^b	115.86±23.58^{ab}	109.85±17.27^{ab}	111.30±40.63^{ab}
	VH/CD	4.16±0.94^b	4.54±1.82^a	4.25±1.07^{ab}	4.38±1.20^{ab}	4.30±2.95^{ab}
Jejunum	VH, um	504.98±50.81	506.92±55.92	507.72±51.80	511.14±54.23	506.89±68.33
	CD, um	100.73±21.13^a	92.21±22.21^b	96.34±30.43^{ab}	95.24±21.78^{ab}	93.18±21.67^b
	VH/CD	4.80±1.59^b	5.29±0.90^a	5.00±1.79^{ab}	5.13±1.28^{ab}	5.23±1.43^a
Ileum	VH, um	518.49±36.85	534.50±62.81	531.09±75.33	537.14±97.02	540.45±49.71
	CD, um	129.72±21.91	135.84±42.42	135.49±17.47	134.77±14.53	136.16±18.01
	VH/CD	4.10±0.66	4.15±1.46	4.12±0.98	4.14±1.23	4.15±0.77

The data in the table are compared in the same row, and different lowercase letters indicate that the difference has reached a significant level (P<0.05); ***CTC:*** *chlortetracycline,* ***FOS:*** *Fructo oligosaccharide,* ***VH:*** *villus height,* ***CD:*** *crypt depth*

Serum Biochemical Indicators

As shown in *Table 3*, There was no significant difference in serum TP, ALB, GLOB, AST, GLU, TC and LDL-C levels among the groups. The ALT level in 100 mg/kg CTC group was significantly higher than in other groups (P<0.05). Adding 100 mg/kg CTC, 2x10^{5}cfu g/kg *B. subtilis* and 200 mg/kg FOS+1x10^{5}cfu g/kg *B. subtilis* can improve the BUN and HDL-C (P<0.05). Significant differences in serum biochemical indicators between groups are shown in *Fig. 2*.

Intestinal Morphology

The intestinal morphology is shown in *Fig. 3*. It can be seen from *Table 4* that in duodenum, the CD of the CTC group was lower than that of the BD group (P<0.05), and the VH/CD were higher (P<0.05), but there is no difference between VH/CD of 100 mg/kg CTC, 2x10^{5}cfu g/kg *B. subtilis* and 200 mg/kg FOS+1x10^{5}cfu g/kg *B. subtilis* groups. In jejunum, the CD of the 100 mg/kg CTC and 200 mg/kg FOS+1x10^{5}cfu g/kg *B. subtilis* groups was lower than that of the BD group (P<0.05), and the VH/CD

were higher (P<0.05). There is no difference between VH, CD and VH/CD among the groups in ileum and there is no difference in the VH of each intestinal tract.

Discussion

Measuring animal growth performance is important because it provides information on the efficiency of feed utilization and overall health of the animal [22]. The ADG and F/G are critical metrics for assessing the growth performance of animals, a high ADG is indicative of strong digestive and absorption capabilities, while a low F/G signifies efficient feed conversion. The better an animal's ability to transform feed into weight gain, the more successful it will be in its growth performance [23]. Antibiotics have a direct impact on the gut microbiome by reducing the population of harmful bacteria and promoting the growth of beneficial bacteria [24]. This can improve gut health and enhance the absorption of nutrients from feed, leading to increased weight gain [25]. Alternatively, using probiotics, prebiotics, and synbiotics can achieve similar results without leaving any residue of antibiotics [26,27]. FOS is non-digestible compounds that stimulate the growth of beneficial bacteria in the gut [28]. Many studies have shown that FOS can improve the growth performance of animals [29]. However, an excessive amount of FOS can also put a strain on the intestines. As a result, FOS is typically used in conjunction with probiotics in animals [30,31]. *B. subtilis* is known for its ability to improve digestive health, increase nutrient absorption, and boost the immune system in animals. This bacterium also has antimicrobial properties, which can help reduce the risk of infection. In studies, the application of *B. subtilis* in livestock has been shown to improve feed efficiency, increase weight gain, and enhance overall performance in chickens, pigs, and ruminants [32-34]. In this experiment, adding 0.2% *B. subtilis* or 0.1% *B. subtilis* + 0.2% FOS increased the final body weight and adding 0.1% *B. subtilis*+0.2% FOS average daily gain, so we think it improved the growth performance of geese.

A higher villus height or shallower crypt depth can contribute to higher digestibility [35,36]. At the same time a higher VH/CD ratio is generally associated with a more efficient small intestine, as there is a greater surface area for nutrient absorption [37]. Conversely, a lower VH/CD ratio is associated with a less efficient small intestine and can be a sign of damage to the intestinal lining [38]. In the experiment, no significant differences were observed in the Ileum indicators among the groups. This is due to the fact that the Ileum is the final segment of the small intestine, where nutrient absorption capacity is limited. Antibiotics can reduce the proliferation of harmful bacteria and promote the differentiation of intestinal epithelial cells, leading to a decrease in the crypt depth and an increase in VH/CD [39]. In our experiment, we observed similar outcomes, however, it is noteworthy that the simultaneous addition of FOS and *B. subtilis* also yielded comparable results, particularly in the jejunum, which has the highest digestion capacity. FOS have been found to enhance the synthesis of polyamines and play a positive role in regulating the growth and development of the small intestine and colonic mucosa. Studies have shown that adding 0.25% FOS to pig feed can increase the villus height and the VH/CD in the proximal small intestine [40]. Additionally, adding 0.4% or 0.6% FOS to pig feed resulted in an increase in villus height and the VH/CD in the jejunum [41]. A study by Howard et al.[42] found that feeding neonatal piglets with FOS led to an increase in the density of cecal epithelial cells and the depth of mucosal crypts in the proximal and distal epithelial cells of the colon. This was attributed to the short-chain fatty acids produced by bacteria metabolizing FOS, which provide energy for intestinal cell proliferation, thus effectively promoting intestinal development. *B. subtilis* has been found to reduce the proliferation of harmful bacteria in the gut, which is likely a contributing factor to the decrease in crypt depth. Several studies have demonstrated that the presence of *B. subtilis* can positively impact the morphological structure of the intestine [43,44]. In this experiment, the combination of *B. subtilis* and FOS was found to have an improved effect on the intestinal morphology of geese.

Serum ALB, GLOB, and TP are important indicators of overall health and nutrition [45]. Despite the absence of significant differences between the groups, we observed an improvement compared with basal diet group. ALT, and AST are commonly used as markers for liver and heart damage or disease [46,47]. Antibiotic-induced liver injury is a known adverse effect of certain antibiotics, and can cause an increase in ALT levels [48]. This increase can be a sign of liver damage, and may be seen in animals receiving certain antibiotics, such as tetracyclines, aminoglycosides, and sulfonamides [49]. The findings of this experiment align with our expectations. The highest levels of both ALT and AST were observed in the group treated with 0.1% CTC. It is worth noting that the level of ALT in the 0.1% CTC group was significantly higher than those seen in any of the other groups. HDL-C, a type of cholesterol commonly referred to as "good" cholesterol. Some antibiotics, such as statins, can lower serum HDL-C level [50], while others may have no significant effect. While the 0.1% CTC group showed elevated levels of HDL-C, it is important to note that this did not diminish the observed liver damage in this group. BUN is a non-protein nitrogenous compound in the blood and is one of the primary products of protein metabolism in the body. Impaired renal function and excessive protein intake can lead to an elevated level of BUN in the blood [51,52]. In this experiment, we found that both CTC and *B. subtilis* can reduce the level of BUN, which indicates that there is more protein or amino acids

degraded into urea in the basal diet group. Both antibiotics and probiotics can reduce the abundance of harmful bacteria, improve intestinal villi, and enhance the ability to absorb protein [19]. Therefore, we believe that adding CTC and *B. subtilis* can both improve the absorption level of protein in geese, and adding 0.1% B. subtilis + 0.2% FOS and 0.1% CTC has a similar effect.

In summary, adding 100 mg/kg *B. subtilis* and $2x10^5$cfu g/kg FOS to the diet as an alternative to antibiotics can significantly improve the growth performance of geese in 14-70 d, improve the intestinal morphology of the jejunum, and prevent liver damage associated with antibiotics.

Availability of Data and Materials

The full dataset and supporting materials are provided within the study for reference. If additional information is desired, please direct inquiries to the corresponding author (H-Y Wang).

Acknowledgements

We would like to express our heartfelt gratitude to the staff at the Zhuanghang Experimental Station, Shanghai, China for their invaluable support and assistance in carrying out this research.

Ethical Approval

Animal experimentation was approved by the Laboratory Animal Ethics Committee of the Shanghai Academy of Agricultural Sciences (SAASPZ0522046).

Funding Support

This study was generously supported by China Agriculture Research System (CARS-42-35), the Climbing Plan of Shanghai Academy of Agricultural Sciences (PG21171), and the Excellent Research Team Program of SAAS (2022-021).

Conflict of Interest

The authors declared that there is no conflict of interest.

Author Contributions

The conception and design of the experiment were contributed by H-Y Wang and D-Q He. G-Q Li and X-Z Wang conducted the data analysis and drafted the manuscript. Yi L, Y-Z Yang, S-M Gong and C Wang were instrumental in data collection. All authors reviewed and approved the final manuscript for publication.

References

1. Castanon JIR: History of the use of antibiotic as growth promoters in European poultry feeds. *Poult Sci*, 86 (11): 2466-2471, 2007. DOI: 10.3382/ps.2007-00249

2. Ahrens P, Andersen LOB, Lilje B, Johannesen TB, Dahl EG, Baig S, Falk L: Changes in the vaginal microbiota following antibiotic treatment for *Mycoplasma genitalium, Chlamydia trachomatis* and bacterial vaginosis. *PloS One*, 15 (7):e0236036, 2020. DOI: 10.1371/journal.pone.0236036

3. Pokhrel A: Antimicrobial (antibiotic) resistance: An evolving threat to medicine. *Ann Med Surg*, 81:104357, 2022. DOI: 10.1016/j.amsu.2022.104357

4. Lavrukhina OI, Amelin VG, Kish, LK, Tretyakov AV, Pen'kov TD: Determination of residual amounts of antibiotics in environmental samples and food products. *J Anal Chem*, 77 (11): 1349-1385, 2022. DOI: 10.1134/S1061934822110077

5. Wang F, Lu Y, Cao J: Dynamics impacts of oxytetracycline on growth performance, intestinal health and antibiotic residue of grouper in exposure and withdrawal treatment. *Ecotox Environ Safe*, 247:114203, 2022. DOI: 10.1016/j.ecoenv.2022.114203

6. Hua Y, Yao Q, Lin J, Li X, Yang Y: Comprehensive survey and health risk assessment of antibiotic residues in freshwater fish in southeast China. *J Food Compos Anal*, 114:104821, 2022. DOI: 10.1016/j.jfca.2022.104821

7. Komura T, Takemoto A, Kosaka H, Suzuki T, Nishikawa Y: Prolonged lifespan, improved perception, and enhanced host defense of *Caenorhabditis elegans* by *Lactococcus cremoris* subsp. *cremoris*. *Microbiol Spectr*, 10 (3):e00454-21, 2022. DOI: 10.1128/spectrum.00454-21

8. Özbek YS, Özmen I: Optimization for coproduction of protease and cellulase from *Bacillus subtilis* M-11 by the Box-Behnken design and their detergentcompatibility. *Braz J Chem Eng*, 37, 49-59, 2020. DOI: 10.1007/s43153-020-00025-x

9. Mutlu A, Trauth S, Ziesack M, Nagler K, Bergeest JP, Rohr K, Bischofs IB: Phenotypic memory in *Bacillus subtilis* links dormancy entry and exit by a spore quantity-quality tradeoff. *Nat Commun*, 9 (1):69, 2018. DOI: 10.1038/s41467-017-02477-1

10. Elegbeleye JA, Buys EM: Molecular characterization and biofilm formation potential of *Bacillus subtilis* and *Bacillus velezensis* in extended shelf-life milk processing line. *J Dairy Sci*, 103 (6): 4991-5002, 2020. DOI: 10.3168/jds.2019-17919

11. Muniyappan M, Baek DH, Kim IH: Effects of dietary supplementation of quillaja saponin or fructooligosaccharide and a mixture of both on the growth performance, nutrient utilisation, faecal microbial and faecal noxious gas emissions in growing pigs. *Ital J Anim Sci*, 21 (1): 1489-1497, 2022. DOI: 10.1080/1828051X.2022.2093656

12. Abd El-Aziz AH, Abo Ghanima MM, Alsanie WF, Gaber A, Alsenosy AE, Easa AA, Moawed SA, Raza SHA, Elfadadny A, Yossef HA, Ghoneem WM, Shukry M, Hendawy AO, Mahrose K: Fructooligosaccharide supplementation boosts growth performance, antioxidant status, and cecal microbiota differently in two rabbit breeds. *Animals*, 12 (12):1528, 2022. DOI: 10.3390/ani12121528

13. Guerreiro I, Serra CR, Oliva-Teles A, Enes P: Gut microbiota of European sea bass *(Dicentrarchus labrax)* is modulated by short-chain fructooligosaccharides and xylooligosaccharides. *Aquacul Int*, 26, 279-288, 2018. DOI: 10.1007/s10499-017-0220-4

14. Bird AR, Vuaran M, Crittenden R, Hayakawa T, Playne MJ, Brown IL, Topping DL: Comparative effects of a high-amylose starch and a fructooligosaccharide on fecal bifidobacteria numbers and short-chain fatty acids in pigs fed Bifidobacterium animalis. *Digest Dis Sci,* 54, 947-954, 2009. DOI: 10.1007/s10620-008-0451-3

15. Porcheron G, Chanteloup NK, Trotereau A, Brée A, Schouler C: Effect of fructooligosaccharide metabolism on chicken colonization by an extra-intestinal pathogenic *Escherichia coli* strain. *PLoS One*, 7 (4):e35475, 2012. DOI: 10.1371/journal.pone.0035475

16. Mirghaed AT, Yarahmadi P, Hosseinifar SH, Tahmasebi D, Gheisvandi N, Ghaedi A: The effects singular or combined administration of fermentable fiber and probiotic on mucosal immune parameters, digestive enzyme activity, gut microbiota and growth performance of Caspian white fish *(Rutilus frisii kutum)* fingerlings. *Fish Shellfish Immun*, 77, 194-199, 2018. DOI: 10.1016/j.fsi.2018.02.007

17. Li WJ: Effects of probiotics, acidifiers and enzymes on growth performance, meat quality, serum biochemistry and antioxidant capacity of Dagu chickens. *Feed Res,* 5, 36-41, 2023.

18. Zhang B, Sui F, Wang B, Wang Y, Li W: Dietary combined supplementation of iron and *Bacillus subtilis* enhances reproductive performance, eggshell quality, nutrient digestibility, antioxidant capacity, and hematopoietic function in breeder geese. *Poult Sci*, 99 (11): 6119-6127, 2020. DOI: 10.1016/j.psj.2020.06.077

19. He T, Long S, Mahfuz S, Wu D, Wang X, Wei X, Piao X: Effects of probiotics as antibiotics substitutes on growth performance, serum biochemical parameters, intestinal morphology, and barrier function of broilers. *Animals*, 9 (11):985, 2019. DOI: 10.3390/ani9110985

20. Dale N: National Research Council (1994) Nutrient Requirements of Poultry. 9th revised ed., National Academy Press, Washington DC, 1994).

21. Xue JJ, Huang XF, Liu ZL, Chen Y, Zhang, YK, Luo Y, Wang C: Effects of citric acid supplementation on growth performance, intestinal

morphology and microbiota, and blood parameters of geese from 1 to 28 days of age. *Poult Sci,* 102 (2):102343, 2023. DOI: 10.1016/j.psj.2022.102343

22. Marx FO, Massuquetto A, Bassi LS, Krabbe E L, Rocha C, Oliveira SG, Maiorka A: Different soybean meal particle sizes on growth performance, nutrient ileal digestibility, digestible energy, and carcass yield of broiler chickens. *Lives Sci,* 247, 104467, 2021. DOI: 10.1016/j.livsci.2021.104467

23. Zhang J, He H, Yuan Y, Wan K, Li L, Liu A: Effects of yeast culture supplementation on growth performance, nutrient digestibility, blood metabolites, and immune response in geese. *Animals*, 12 (10):1270, 2022. DOI: 10.3390/ani12101270

24. Cremonesi P, Biscarini F, Castiglioni B, Sgoifo CA, Compiani R, Moroni P: Gut microbiome modifications over time when removing in-feed antibiotics from the prophylaxis of post-weaning diarrhea in piglets. *PloS One*, 17 (3):e0262199, 2022. DOI: 10.1371/journal.pone.0262199

25. Wang Y, Wang Y, Lin X, Gou Z, Fan Q, Ye J, Jiang S: Potential effects of acidifier and amylase as substitutes for antibiotic on the growth performance, nutrient digestion and gut microbiota in yellow-feathered broilers. *Animals*, 10 (10):1858, 2020. DOI: 10.3390/ani10101858

26. Wang B, Zhou Y, Tang L, Zeng Z, Gong L, Wu Y, Li WF: Effects of Bacillus amyloliquefaciens instead of antibiotics on growth performance, intestinal health, and intestinal microbiota of broilers. *Front Vet Sci*, 8:679368, 2021. DOI: 10.3389/fvets.2021.679368

27. Wang LM, Mandell IB, Bohrer BM: Effects of feeding essential oils and benzoic acid to replace antibiotics on finishing beef cattle growth, carcass characteristics, and sensory attributes. *Appl Anim Sci*, 36 (2): 145-156, 2020. DOI: 10.15232/aas.2019-01908

28. Yan H, Zhou P, Zhang Y, Zhang Z, Liu J, Zhang H: Short-chain fructo-oligosaccharides alleviates oxidized oil-induced intestinal dysfunction in piglets associated with the modulation of gut microbiota. *J Funct Foods,* 64:103661, 2020. DOI: 10.1016/j.jff.2019.103661

28. Liu ZY, Yang HL, Yan YY, Seerengaraj V, Zhang CX, Ye JD, Sun YZ: Supplementation of tributyrin, alone and in combination with fructo-oligosaccharide in high soybean meal diets for shrimp *(Litopenaeus vannamei)*: Effects on growth, innate immunity and intestinal morphology. *Aquacul Nutr*, 27 (2): 592-603, 2021. DOI: 10.1111/anu.13209

29. Zhang CN, Li XF, Xu WN, Zhang DD, Lu KL, Wang LN, Liu WB: Combined effects of dietary fructooligosaccharide and *Bacillus licheniformis* on growth performance, body composition, intestinal enzymes activities and gut histology of triangular bream *(Megalobrama terminalis). Aquacul Nutr*, 21 (5): 755-766, 2015. DOI: 10.1111/anu.12200

30. Bomhof MR, Saha DC, Reid DT, Paul HA, Reimer RA: Combined effects of oligofructose and Bifidobacterium animalis on gut microbiota and glycemia in obese rats. *Obesity*, 22 (3): 763-771, 2014. DOI: 10.1002/oby.20632

31. Sen S, Ingale SL, Kim JS, Kim KH, Kim YW, Khong C, Chae BJ: Effect of supplementation of *Bacillus subtilis* LS 1-2 grown on citrus-juice waste and corn-soybean meal substrate on growth performance, nutrient retention, caecal microbiology and small intestinal morphology of broilers. *Asian-Australas J Anim Sci*, 24 (8): 1120-1127, 2011. DOI: 10.5713/ajas.2011.10443

32. Hossain MM, Begum M, Kim IH: Effect of *Bacillus subtilis, Clostridium butyricum* and Lactobacillus acidophilus endospores on growth performance, nutrient digestibility, meat quality, relative organ weight, microbial shedding and excreta noxious gas emission in broilers. *Vet Med*, 60 (2): 77-86, 2015. DOI: 10.17221/7981-VETMED

33. Lee SH, Ingale SL, Kim JS, Kim KH, Lokhande A, Kim EK, Chae BJ: Effects of dietary supplementation with *Bacillus subtilis* LS 1-2 fermentation biomass on growth performance, nutrient digestibility, cecal microbiota and intestinal morphology of weanling pig. *Anim Feed Sci Technol*, 188, 102-110, 2014. DOI: 10.1016/j.anifeedsci.2013.12.001

34. Fukumori R, DOI K, Mochizuki T, Oikawa S, Gondaira S, Iwasaki T, Izumi K: Sodium butyrate administration modulates the ruminal villus height, inflammation-related gene expression, and plasma hormones concentration in dry cows fed a high-fiber diet. *Anim Sci J*, 93 (1):e13791, 2022. DOI: 10.1111/asj.13791

35. Chwen LT, Foo HL, Thanh NT, Choe DW: Growth performance, plasma Fatty acids, villous height and crypt depth of preweaning piglets fed with medium chain triacylglycerol. *Asian-Australas J Anim Sci,* 26 (5): 700-704, 2013. DOI: 10.5713/ajas.2012.12561

36. Gurung M, Rosa F, Yelvington B, Terry N, Read QD, Piccolo BD, Yeruva L: Evaluation of a plant-based infant formula containing almonds and buckwheat on gut microbiota composition, intestine morphology, metabolic and immune markers in a neonatal piglet model. *Nutrients*, 15 (2):383, 2023. DOI: 10.3390/nu15020383

37. Xu Y, Zhang B, Zhao R, Gao K, Liang S, Wu Y, Tang J: Effect of riboflavin deficiency on intestinal morphology, jejunum mucosa proteomics, and cecal microbiota of Pekin ducks. *Anim Nutr*, 12, 215-226, 2023. DOI: 10.1016/j.aninu.2022.09.013

38. Zhang Q, Li J, Yi X, Li Z, Liang S, Fang Z, Che L: Rhodotorula benthica culture as an alternative to antibiotics improves growth performance by improving nutrients digestibility and intestinal morphology, and modulating gut microbiota of weaned piglets. *Front Microbiol*, 13:964651, 2022. DOI: 10.3389/fmicb.2022.964531

40. Shim S: Effects of prebiotics, probiotics and synbiotics in the diet of young pigs. *PhD Thesis,* Wageningen Univ Res, 28238877, 2005.

41. Xu ZR, Zou XT, Hu CH, Xia MS, Zhan XA, Wang MQ: Effects of dietary fructooligosaccharide on digestive enzyme activities, intestinal microflora and morphology of growing pigs. *Asian-Australas J Anim Sci*, 15 (12): 1784-1789, 2022. DOI: 10.5713/ajas.2002.1784

42. Howard MD, Gordon DT, Pace LW, Garleb KA, Kerley MS: Effects of dietary supplementation with fructooligosaccharides on colonic microbiota populations and epithelial cell proliferation in neonatal pigs. *J Pediatr Gastr Nutr*, 21 (3): 297-303, 1995. DOI: 10.1097/00005176-199510000-00007

43. Xing Y, Wang S, Fan J, Oso AO, Kim SW, Xiao D, Zhang B: Effects of dietary supplementation with lysine-yielding Bacillus subtilis on gut morphology, cecal microflora, and intestinal immune response of Linwu ducks. *J Anim Sci*, 93 (7): 3449-3457, 2015. DOI: 10.2527/jas.2014-8090

44. Al-Fataftah AR, Abdelqader A: Effects of dietary Bacillus subtilis on heat-stressed broilers performance, intestinal morphology and microflora composition. *Anim Feed Sci Technol*, 198, 279-285, 2014. DOI: 10.1016/j.anifeedsci.2014.10.012

45. Ölmez M, Şahin T, Özlem Ö, Öğün M, Yörük MA, Dalğa S: Effect of probiotic mixture supplementation to drinking water on the growth performance, carcass parameters and serum biochemical parameters in native Turkish geese. *Kafkas Univ Vet Fak Derg*, 28 (1): 131-138, 2022. DOI: 10.9775/kvfd.2021.26633

46. Zhu CW, Liu JC, Wang J, Wang H, Liu YL, Li YG, Zhu L, Ding WM, Qiu YW, Yang YF, Li J, Huang R, Wu C: Liver inflammation activity in autoimmune hepatitis patients with normal ALT and IgG levels. *J Hepatol*, 77, S326-S327, 2022. DOI: 10.1016/S0168-8278(22)01017-0

47. Kogan M, Klempfner R, Lotan D, Wasserstrum Y, Goldenberg I, Segal G: Low ALT blood levels are associated with lower baseline fitness amongst participants of a cardiac rehabilitation program. *J Exerc Sci Fit*, 16 (1): 1-4, 2018. DOI: 10.1016/j.jesf.2017.11.002

48. Liu P, Yang M, Jia CX, Zhang QJ, Qu JQ: Effect of enrofloxacin on transaminase activity in serum and liver of Siberian sturgeon. *Fish Sci*, 2, 273-279, 2022. DOI: 10.16378/j.cnki.1003-1111.20090

49. Haque MI, Ahmad N, Miah MA: Comparative analysis of body weight and serum biochemistry in broilers supplemented with some selected probiotics and antibiotic growth promoters. *J Adv Vet Anim Res*, 4 (3): 288-294, 2017. DOI: 10.5455/javar.2017.d226

50. Mori T, Tanno Y, Kasakura S, Yoshioka K, Nakai N: Influence of statins on LDL-and HDL-cholesterol and plasma fatty acids in elderly Japanese ischemic stroke patients. *Austin J Cerebrovasc Dis Stroke*, 4 (3): 1062-1064, 2017. DOI: 10.26420/AUSTINJCEREBROVASCDISSTROKE.2017.1062

51. Hou H, Li L: Effects of continuous renal replacement therapy on Apache-II score, creatinine, and urea nitrogen levels in patients with acute kidney injury. *Pak J Med Sci*, 39 (1):6591, 2023. DOI: 10.12669/pjms.39.1.6591

52. Yu D, Zhu W, Hang S: Effects of low-protein diet on the intestinal morphology, digestive enzyme activity, blood urea nitrogen, and gut microbiota and metabolites in weaned pigs. *Arch Anim Nutr*, 73 (4): 287-305, 2019. DOI: 10.1080/1745039X.2019.1614849