Hacettepe Journal of Mathematics & Statistics

Hacet. J. Math. Stat.
Volume 54 (5) (2025), 1826 – 1838
DOI : 10.15672/hujms.1579443

Research Article

On the numerical radii of certain operator forms

Fuad Kittaneh*[1,2], Hamid Reza Moradi[3], Mohammad Sababheh[3]

[1] *Department of Mathematics, The University of Jordan, Amman, Jordan*
[2] *Department of Mathematics, Korea University, Seoul, 02841, South Korea*
[3] *Department of Mathematics, Mashhad Branch, Islamic Azad University, Mashhad, Iran*
[4] *Department of Basic Sciences, Princess Sumaya University for Technology, Amman, Jordan*

Abstract

The main goal of this paper is to present new bounds for the norms and numerical radii of certain Hilbert space operators, which involve the sums of operators' products. The obtained result will be utilized to obtain some equivalent conditions regarding certain operator identities, to find upper bounds for the sum of two operators, to obtain certain refinements of celebrated numerical radius bounds, and to present a new reverse for the triangle inequality when positive operators are treated.

Mathematics Subject Classification (2020). Primary 47A12, 47A30, Secondary 15A18, 47A63

Keywords. numerical radius, operator norm, spectral radius, operator inequality

1. Introduction

In the sequel, $\mathbb{H}$ is a complex Hilbert space, endowed with inner product $\langle\cdot,\cdot\rangle$, and induced norm $\|\cdot\|$. The set of all bounded linear operators from $\mathbb{H}$ to $\mathbb{H}$ will be denoted by $\mathbb{B}(\mathbb{H})$, with zero element O and identity I. Upper case letters through this paper will be used to denote elements of $\mathbb{B}(\mathbb{H})$.

Associated with $A \in \mathbb{B}(\mathbb{H})$, many scalar values have received the attention of specialists through the literature. Among those quantities, the operator norm, the numerical radius, and the spectral radius have received researchers' attention.

These three quantities are defined, respectively, for an arbitrary $A \in \mathbb{B}(\mathbb{H})$ as follows

$$\|A\| = \sup_{\|x\|=1} \|Ax\|, \omega(A) = \sup_{\|x\|=1} |\langle Ax, x\rangle| \text{ and } r(A) = \sup\{|\lambda| : \lambda \in \sigma(A)\},$$

where $\sigma(A)$ denotes the spectrum of A, defined as $\sigma(A) = \{\lambda \in \mathbb{C} : A - \lambda I \text{ is not invertible}\}$.

It is well known that $\|A\| = \sup_{\|x\|=\|y\|=1} |\langle Ax, y\rangle|$, which makes the relation $\omega(A) \leq \|A\|$ clear, for any $A \in \mathbb{B}(\mathbb{H})$.

*Corresponding Author.
Email addresses: fkittr@ju.edu.jo (F. Kittaneh), hrmoradi@mshdiau.ac.ir (H. R. Moradi), sababheh@yahoo.com (M. Sababheh), sababheh@psut.edu.jo (M. Sababheh)
Received: 05.11.2024; Accepted: 11.02.2025

Among those interesting investigations in the literature is the investigation of possible relations that govern the quantities $\|\cdot\|, \omega(\cdot)$ and $r(\cdot)$. It is well established that for any $A \in \mathbb{B}(\mathbb{H})$, one has [19, Theorem 1.3-1 and Theorem 1.4-4]

$$r(A) \leq \omega(A) \leq \|A\|. \tag{1.1}$$

However, discussing sharper forms of (1.1) has been an exciting topic. This interest has received considerable attention, and hundreds of research papers have been published. We refer the interested reader to [2–8, 10, 12–15, 18, 20, 22, 23, 30, 31, 33–38] and the references therein, as a sample of such work.

Letting A^* denote the adjoint of $A \in \mathbb{B}(\mathbb{H})$, we say that A is normal if $A^*A = AA^*$. The class of normal operators fulfills sharper bounds than general operators. For example, both inequalities in (1.1) become equalities [19, Theorem 1.4-2].

While simple forms are desired to describe such relations, the literature has witnessed more elaborated forms that had some influence on the advancement of this field. In particular, operator matrices have played a vital role in understanding some basic properties of the numerical radius and its relation with the operator norm and spectral radius. We recall here that if $A, B, C, D \in \mathbb{B}(\mathbb{H})$, then the operator matrix $\begin{bmatrix} A & B \\ C & D \end{bmatrix} \in \mathbb{B}(\mathbb{H} \oplus \mathbb{H})$, where $\oplus$ denotes the direct sum. We refer the reader to [1, 9, 14, 29, 32] as a sample of how operator matrices influenced and enriched the study of the above interest.

In this paper, we discuss possible bounds in this context, where sums of products of certain operators are treated. We refer the reader to [21], where the quantity $\omega(AXB^* + BYA^*)$ has been studied, for $A, B, X, Y \in \mathbb{B}(\mathbb{H})$. It is interesting that the upper bound for this quantity is found in terms of an operator matrix.

Given $A_1, A_2, B_1, B_2 \in \mathbb{B}(\mathbb{H})$, we will discuss possible bounds for the operator norm and the numerical radius of the operator $A_1B_1 + A_2B_2$. The obtained bounds are used to obtain certain bounds for the norm of the sum of two operators and the norm of an operator's Cartesian parts. Here, we recall that if $A \in \mathbb{B}(\mathbb{H})$, the real and imaginary parts of A are defined, respectively, as

$$\Re A = \frac{A + A^*}{2} \text{ and } \Im A = \frac{A - A^*}{2\mathrm{i}}.$$

The real and imaginary parts have strong relations to the numerical radius, as one can verify that

$$\omega(A) = \sup_{\theta \in \mathbb{R}} \left\| \Re\left(e^{\mathrm{i}\theta} A\right) \right\| = \sup_{\theta \in \mathbb{R}} \left\| \Im\left(e^{\mathrm{i}\theta} A\right) \right\|.$$

We will need some lemmas from the literature to obtain our results, as follows.
For the first lemma, we recall that an operator $A \in \mathbb{B}(\mathbb{H})$ is said to be positive if $\langle Ax, x\rangle \geq 0$ for all $x \in \mathbb{H}$.

Lemma 1.1. [25] Let $A, B \in \mathbb{B}(\mathbb{H})$ be two positive operators. Then

$$\|A + B\| \leq \max\{\|A\|, \|B\|\} + \left\|A^{\frac{1}{2}} B^{\frac{1}{2}}\right\|.$$

The following is an arithmetic-geometric mean inequality; see [11].

Lemma 1.2. *Let $A, B \in \mathbb{B}(\mathbb{H})$. Then*

$$\|AB^*\| \leq \frac{1}{2} \left\| |A|^2 + |B|^2 \right\|,$$

where $|X|^2 = X^*X$ for $X \in \mathbb{B}(\mathbb{H})$.

Lemma 1.3. [16, p. 58] Let $A = U|A|$ be the polar decomposition of A. Then for any $q > 0$,

$$U|A|^q U^* = |A^*|^q.$$

Lemma 1.4. [24] Let $\mathbb{H}_1, \mathbb{H}_2, \dots, \mathbb{H}_n$ be Hilbert spaces, and let $\mathbb{T} = [T_{ij}]$ be an $n \times n$ operator matrix with $T_{ij} \in \mathbb{B}(\mathbb{H}_i, \mathbb{H}_j)$. Then

$$\|\mathbb{T}\| \leq \|[\|T_{ij}\|]\|,$$

2. Main results

We present our results in two subsections, treating related bounds for the operator norm and the numerical radius.

2.1. Upper bounds for the operator norm

The first result gives an elaborated upper bound for $\|A_1B_1 + A_2B_2\|$. Some applications will follow.

Theorem 2.1. *Let $A_1, A_2, B_1, B_2 \in \mathbb{B}(\mathbb{H})$. Then*

$$\|A_1B_1 + A_2B_2\| \leq \frac{1}{2}\sqrt{\|A_1A_1^* + A_2A_2^*\|\,\|B_1^*B_1 + B_2^*B_2\|} + \frac{1}{2}\omega^{\frac{1}{2}}\left(\begin{bmatrix} A_1^*A_1B_1B_1^* + A_1^*A_2B_2B_1^* & A_1^*A_1B_1B_2^* + A_1^*A_2B_2B_2^* \\ A_2^*A_1B_1B_1^* + A_2^*A_2B_2B_1^* & A_2^*A_1B_1B_2^* + A_2^*A_2B_2B_2^* \end{bmatrix}\right).$$

Proof. For any $\mathbf{A}, \mathbf{B} \in \mathbb{B}(\mathbb{H})$, Lemma 1.2 implies

$$\|\mathbf{A}\mathbf{B}^*\| \leq \frac{1}{2}\left\||\mathbf{A}|^2 + |\mathbf{B}|^2\right\|. \tag{2.1}$$

For nonzero operators $\mathbf{A}$ and $\mathbf{B}$, if we replace $\mathbf{A}$ and $\mathbf{B}$ by $\sqrt{\frac{\|\mathbf{B}\|}{\|\mathbf{A}\|}}\mathbf{A}$ and $\sqrt{\frac{\|\mathbf{A}\|}{\|\mathbf{B}\|}}\mathbf{B}$, in (2.1), we infer that

$$\begin{aligned}
\|\mathbf{A}\mathbf{B}^*\| &\leq \frac{1}{2}\left\|\frac{\|\mathbf{B}\|}{\|\mathbf{A}\|}|\mathbf{A}|^2 + \frac{\|\mathbf{A}\|}{\|\mathbf{B}\|}|\mathbf{B}|^2\right\| \\
&\leq \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \||\mathbf{A}|\,|\mathbf{B}|\|\right) \quad \text{(by Lemma 1.1)} \\
&= \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \left\||\mathbf{A}|\,|\mathbf{B}|^2\,|\mathbf{A}|\right\|^{\frac{1}{2}}\right) \\
&= \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \sqrt{r\left(|\mathbf{A}|\,|\mathbf{B}|^2\,|\mathbf{A}|\right)}\right) \\
&= \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \sqrt{r\left(|\mathbf{A}|^2|\mathbf{B}|^2\right)}\right) \\
&\leq \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \omega^{\frac{1}{2}}\left(|\mathbf{A}|^2|\mathbf{B}|^2\right)\right),
\end{aligned}$$

where we have used the fact that $\|X\|^2 = \|X^*\| = \||X|^2\|$ to obtain the first equality, that $r(X) = \|X\|$ when X is Hermitian to obtain the second equality, that $r(\cdot)$ is commutative to obtain the third equality and that $r(\cdot) \leq \omega(\cdot)$ to obtain the last inequality. Thus, we have shown that for all $\mathbf{A}, \mathbf{B} \in \mathbb{B}(\mathbb{H})$,

$$\|\mathbf{A}\mathbf{B}^*\| \leq \frac{1}{2}\left(\|\mathbf{A}\|\,\|\mathbf{B}\| + \omega^{\frac{1}{2}}\left(|\mathbf{A}|^2|\mathbf{B}|^2\right)\right). \tag{2.2}$$

Now for the given A_i, B_i, let $\mathbf{A} = \begin{bmatrix} A_1 & A_2 \\ O & O \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} B_1^* & B_2^* \\ O & O \end{bmatrix}$. Simple calculations reveal that

$$\|\mathbf{A}\mathbf{B}^*\| = \left\|\begin{bmatrix} A_1B_1 + A_2B_2 & O \\ O & O \end{bmatrix}\right\| = \|A_1B_1 + A_2B_2\|,$$

$$\|\mathbf{A}\| = \|\mathbf{A}\mathbf{A}^*\|^{\frac{1}{2}} = \left\|\begin{bmatrix} A_1A_1^* + A_2A_2^* & O \\ O & O \end{bmatrix}\right\|^{\frac{1}{2}} = \|A_1A_1^* + A_2A_2^*\|^{\frac{1}{2}},$$

$$\|\mathbf{B}\| = \|\mathbf{B}\mathbf{B}^*\|^{\frac{1}{2}} = \left\|\begin{bmatrix} B_1^*B_1 + B_2^*B_2 & O \\ O & O \end{bmatrix}\right\|^{\frac{1}{2}} = \|B_1^*B_1 + B_2^*B_2\|^{\frac{1}{2}},$$

and

$$\begin{aligned}&\omega^{\frac{1}{2}}\left(|\mathbf{A}|^2|\mathbf{B}|^2\right)\\ &= \omega^{\frac{1}{2}}\left(\begin{bmatrix} A_1^*A_1B_1B_1^* + A_1^*A_2B_2B_1^* & A_1^*A_1B_1B_2^* + A_1^*A_2B_2B_2^* \\ A_2^*A_1B_1B_1^* + A_2^*A_2B_2B_1^* & A_2^*A_1B_1B_2^* + A_2^*A_2B_2B_2^* \end{bmatrix}\right).\end{aligned}$$

We obtain the desired result by taking into account (2.2). This completes the proof for the nonzero operators case. The zero case is trivial. □

Remark 2.2. For any $A_1, A_2, B_1, B_2 \in \mathbb{B}(\mathbb{H})$, we have

$$\begin{aligned}&\omega\left(\begin{bmatrix} A_1^*A_1B_1B_1^* + A_1^*A_2B_2B_1^* & A_1^*A_1B_1B_2^* + A_1^*A_2B_2B_2^* \\ A_2^*A_1B_1B_1^* + A_2^*A_2B_2B_1^* & A_2^*A_1B_1B_2^* + A_2^*A_2B_2B_2^* \end{bmatrix}\right)\\ &= \omega\left(|\mathbf{A}|^{\mathbf{2}}|\mathbf{B}|^{\mathbf{2}}\right) \leq \left\||\mathbf{A}|^{\mathbf{2}}|\mathbf{B}|^{\mathbf{2}}\right\| \leq \|\mathbf{A}\|^2 \ \|\mathbf{B}\|^2\\ &= \|A_1A_1^* + A_2A_2^*\| \, \|B_1^*B_1 + B_2^*B_2\|.\end{aligned}$$

Consequently, Theorem 2.1 implies

$$\|A_1B_1 + A_2B_2\| \leq \sqrt{\|A_1A_1^* + A_2A_2^*\| \, \|B_1^*B_1 + B_2^*B_2\|}.$$

Following the proof of Theorem 2.1, we may state the following result, which gives multiple equivalent statements that involve certain identities, which are inequalities in general. One can refer to [26] for a similar discussion.

Theorem 2.3. *Let $A, B \in \mathbb{B}(\mathbb{H})$. Then the following statements are equivalent:*

(i) $\|AB\| = \|A\| \, \|B\|$;
(ii) $\omega^{\frac{1}{2}}\left(|A|^2|B^*|^2\right) = \|A\| \, \|B\|$;
(iii) $\left\||A|^2 + |B^*|^2\right\| = \|A\|^2 + \|B\|^2$;
(iv) $\left\||A|^2|B^*|^2\right\| = \|A\|^2\|B\|^2$;
(v) $\sqrt{r\left(|A|^2|B^*|^2\right)} = \|A\| \, \|B\|$.

Proof. If $A = O$ or $B = O$, the result follows immediately. So, without loss of generality, we may assume that A and B are both nonzero.
$(i) \Rightarrow (ii)$ If $\|AB\| = \|A\| \, \|B\|$, then by (2.2), we get

$$\|A\| \, \|B\| \leq \omega^{\frac{1}{2}}\left(|A|^2|B^*|^2\right). \tag{2.3}$$

On the other hand, we know that

$$\omega^{\frac{1}{2}}\left(|A|^2|B^*|^2\right) \leq \left\||A|^2|B^*|^2\right\|^{\frac{1}{2}} \leq \left\||A|^2\right\|^{\frac{1}{2}}\left\||B^*|^2\right\|^{\frac{1}{2}} = \| \, |A| \, \| \, \| \, |B^*| \, \| = \|A\| \, \|B\|. \tag{2.4}$$

Inequalities (2.3) and (2.4) imply $\omega^{\frac{1}{2}}\left(|A|^2|B^*|^2\right) = \|A\| \, \|B\|$.

$(ii) \Rightarrow (i)$ Assume that $\omega^{\frac{1}{2}}\left(|A|^2|B^*|^2\right) = \|A\| \, \|B\|$. Then then there exists a sequence of unit vectors $\{x_n\}$ such that $\|A\|^2\|B\|^2 = \lim\limits_{n\to\infty} |\langle A^*ABB^*x_n, x_n\rangle|$, and so

$$\|A\|^2\|B\|^2 = \lim_{n\to\infty} |\langle ABB^*x_n, Ax_n\rangle| \leq \|AB\| \limsup_{n\to\infty} \|Ax_n\| \, \|B^*x_n\| \leq \|AB\| \, \|A\| \, \|B\|.$$

Hence, $\|A\| \, \|B\| \leq \|AB\|$. But the inequality $\|AB\| \leq \|A\| \, \|B\|$ is always true. These two inequalities imply $\|AB\| = \|A\| \, \|B\|$.

$(i) \Leftrightarrow (iii)$ See [26].
$(i) \Leftrightarrow (iv)$ See [26].
$(i) \Leftrightarrow (v)$ This follows from the following observation:

$$\|AB\| = \| \, |A| \, |B^*| \, \| = \|(|A| \, |B^*|) \, (|A| \, |B^*|)^*\|^{\frac{1}{2}} = \sqrt{r\left(|A|^2 |B^*|^2\right)}.$$

□

Now, as an application of Theorem 2.1, we present the following upper bound for the norm of the sum of two operators.

Corollary 2.4. *Let $S, T \in \mathbb{B}(\mathbb{H})$ be invertible. Then for any $0 \le t \le 1$,*

$$\begin{aligned}\|S+T\| &\le \frac{1}{2}\sqrt{\left\||S|^{2t} + |T|^{2(1-t)}\right\| \left\||S^*|^{2(1-t)} + |T^*|^{2t}\right\|} \\ &\quad + \frac{1}{2}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |S|^2 + |S|^{-t}S^*T|S|^t & |S|^{2-t}|T|^{1-t} + |S|^{-t}S^*T|T|^{1-t} \\ |T|^{1+t}|S|^t + |T|^{t-1}T^*S|S|^t & |T|^2 + |T|^{t-1}T^*S|T|^{1-t} \end{bmatrix}\right) \\ &\le \sqrt{\left\||S|^{2t} + |T|^{2(1-t)}\right\| \left\||S^*|^{2(1-t)} + |T^*|^{2t}\right\|}.\end{aligned}$$

In particular,

$$\begin{aligned}\|S+T\| &\le \frac{1}{2}\sqrt{\| \, |S| + |T| \, \| \, \| \, |S^*| + |T^*| \, \|} \\ &\quad + \frac{1}{2}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |S|^2 + |S|^{-\frac{1}{2}}S^*T|S|^{\frac{1}{2}} & |S|^{\frac{3}{2}}|T|^{\frac{1}{2}} + |S|^{-\frac{1}{2}}S^*T|T|^{\frac{1}{2}} \\ |T|^{\frac{3}{2}}|S|^{\frac{1}{2}} + |T|^{-\frac{1}{2}}T^*S|S|^{\frac{1}{2}} & |T|^2 + |T|^{-\frac{1}{2}}T^*S|T|^{\frac{1}{2}} \end{bmatrix}\right) \\ &\le \sqrt{\| \, |S| + |T| \, \| \, \| \, |S^*| + |T^*| \, \|}.\end{aligned}$$

Proof. Let $S = U\,|S|$ and $T = V\,|T|$ be the polar decompositions of S and T, respectively. Let $A_1 = U|S|^{1-t}$, $B_1 = |S|^t$, $A_2 = V|T|^t$, and $B_2 = |T|^{1-t}$. Then we observe that

$$\begin{aligned}A_1A_1^* + A_2A_2^* &= U|S|^{1-t}|S|^{1-t}U^* + V|T|^t|T|^tV^* \\ &= U|S|^{2(1-t)}U^* + V|T|^{2t}V^* \\ &= |S^*|^{2(1-t)} + |T^*|^{2t} \quad \text{(by Lemma 1.3)},\end{aligned}$$

$$B_1^*B_1 + B_2^*B_2 = |S|^t|S|^t + |T|^{1-t}|T|^{1-t} = |S|^{2t} + |T|^{2(1-t)},$$

$$\begin{aligned}&A_1^*A_1B_1B_1^* + A_1^*A_2B_2B_1^* \\ &= |S|^{1-t}U^*U|S|^{1-t}|S|^t|S|^t + |S|^{1-t}U^*V|T|^t|T|^{1-t}|S|^t \\ &= |S|^{1-t}U^*U|S|^{1-t}|S|^{2t} + |S|^{1-t}U^*V\,|T|\,|S|^t \\ &= |S|^2 + |S|^{1-t}U^*V\,|T|\,|S|^t \quad \text{(since } U^*U \text{ is a projection)} \\ &= |S|^2 + |S|^{-t}\,|S|\,U^*V\,|T|\,|S|^t \\ &= |S|^2 + |S|^{-t}S^*T|S|^t \quad \text{(since } S^* = |S|\,U^* \text{ and } T = V\,|T|),\end{aligned}$$

$$\begin{aligned}&A_1^*A_1B_1B_2^* + A_1^*A_2B_2B_2^* \\ &= |S|^{1-t}U^*U|S|^{1-t}|S|^t|T|^{1-t} + |S|^{1-t}U^*V|T|^t|T|^{1-t}|T|^{1-t} \\ &= |S|^{2-t}|T|^{1-t} + |S|^{1-t}U^*V\,|T|\,|T|^{1-t} \quad \text{(since } U^*U \text{ is projection)} \\ &= |S|^{2-t}|T|^{1-t} + |S|^{-t}\,|S|\,U^*V\,|T|\,|T|^{1-t} \\ &= |S|^{2-t}|T|^{1-t} + |S|^{-t}S^*T|T|^{1-t} \quad \text{(since } S^* = |S|\,U^* \text{ and } T = V\,|T|),\end{aligned}$$

$$\begin{aligned}
&A_2^*A_1B_1B_1^* + A_2^*A_2B_2B_1^*\\
&= |T|^tV^*U|S|^{1-t}|S|^t|S|^t + |T|^tV^*V|T|^t|T|^{1-t}|S|^t\\
&= |T|^tV^*U\,|S|\,|S|^t + |T|^{1+t}|S|^t \quad \text{(since } V^*V \text{ is projection)}\\
&= |T|^{t-1}\,|T|\,V^*U\,|S|\,|S|^t + |T|^{1+t}|S|^t\\
&= |T|^{t-1}T^*S|S|^t + |T|^{1+t}|S|^t \quad \text{(since } T^* = |T|\,V^* \text{ and } S = U\,|S|),
\end{aligned}$$

and

$$\begin{aligned}
&A_2^*A_1B_1B_2^* + A_2^*A_2B_2B_2^*\\
&= |T|^tV^*U|S|^{1-t}|S|^t|T|^{1-t} + |T|^tV^*V|T|^t|T|^{1-t}|T|^{1-t}\\
&= |T|^tV^*U\,|S|\,|T|^{1-t} + |T|^2 \quad \text{(since } V^*V \text{ is a projection)}\\
&= |T|^{t-1}\,|T|\,V^*U\,|S|\,|T|^{1-t} + |T|^2\\
&= |T|^{t-1}T^*S|T|^{1-t} + |T|^2 \quad \text{(since } T^* = |T|\,V^* \text{ and } S = U\,|S|).
\end{aligned}$$

The result follows by Theorem 2.1, noting Remark 2.2. □

Remark 2.5. Let $T \in \mathbb{B}(\mathbb{H})$.

(i) Replacing S by T^* in Corollary 2.4 yields

$$\begin{aligned}
\|\Re T\| &\le \frac{1}{4}\left\||T|^{2(1-t)} + |T^*|^{2t}\right\|\\
&\quad + \frac{1}{4}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |T^*|^2 + |T^*|^{-t}T^2|T^*|^t & |T^*|^{2-t}|T|^{1-t} + |T^*|^{-t}T^2|T|^{1-t}\\ |T|^{1+t}|T^*|^t + |T|^{t-1}(T^*)^2|T^*|^t & |T|^2 + |T|^{t-1}(T^*)^2|T|^{1-t}\end{bmatrix}\right)\\
&\le \frac{1}{2}\left\||T|^{2(1-t)} + |T^*|^{2t}\right\|.
\end{aligned}$$

(ii) Replacing T by T^* and S by $-T$ in Corollary 2.4 yield

$$\begin{aligned}
\|\Im T\| &\le \frac{1}{4}\left\||T^*|^{2(1-t)} + |T|^{2t}\right\|\\
&\quad + \frac{1}{4}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |T|^2 - |T|^{-t}(T^*)^2|T|^t & |T|^{2-t}|T^*|^{1-t} - |T|^{-t}(T^*)^2|T^*|^{1-t}\\ |T^*|^{1+t}|T|^t - |T^*|^{t-1}T^2|T|^t & |T^*|^2 - |T^*|^{t-1}T^2|T^*|^{1-t}\end{bmatrix}\right)\\
&\le \frac{1}{2}\left\||T^*|^{2(1-t)} + |T|^{2t}\right\|.
\end{aligned}$$

In particular,

(iii)

$$\begin{aligned}
\|\Re T\| &\le \frac{1}{4}\,\|\,|T| + |T^*|\,\|\\
&\quad + \frac{1}{4}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |T^*|^2 + |T^*|^{-\frac{1}{2}}T^2|T^*|^{\frac{1}{2}} & |T^*|^{\frac{3}{2}}|T|^{\frac{1}{2}} + |T^*|^{-\frac{1}{2}}T^2|T|^{\frac{1}{2}}\\ |T|^{\frac{3}{2}}|T^*|^{\frac{1}{2}} + |T|^{-\frac{1}{2}}(T^*)^2|T^*|^{\frac{1}{2}} & |T|^2 + |T|^{-\frac{1}{2}}(T^*)^2|T|^{\frac{1}{2}}\end{bmatrix}\right)\\
&\le \frac{1}{2}\,\|\,|T| + |T^*|\,\|.
\end{aligned}$$

(iv)

$$\begin{aligned}
\|\Im T\| &\le \frac{1}{4}\,\|\,|T| + |T^*|\,\|\\
&\quad + \frac{1}{4}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |T|^2 - |T|^{-\frac{1}{2}}(T^*)^2|T|^{\frac{1}{2}} & |T|^{\frac{3}{2}}|T^*|^{\frac{1}{2}} - |T|^{-\frac{1}{2}}(T^*)^2|T^*|^{\frac{1}{2}}\\ |T^*|^{\frac{3}{2}}|T|^{\frac{1}{2}} - |T^*|^{-\frac{1}{2}}T^2|T|^{\frac{1}{2}} & |T^*|^2 - |T^*|^{-\frac{1}{2}}T^2|T^*|^{\frac{1}{2}}\end{bmatrix}\right)\\
&\le \frac{1}{2}\,\|\,|T| + |T^*|\,\|.
\end{aligned}$$

Remark 2.6. If S, T are normal operators, then Corollary 2.4 ensures that

$$\begin{aligned}\|S+T\| &\leq \frac{1}{2}\| \, |S|+|T| \, \| \\ &\quad + \frac{1}{2}\omega^{\frac{1}{2}}\left(\begin{bmatrix} |S|^2+|S|^{-\frac{1}{2}}S^*T|S|^{\frac{1}{2}} & |S|^{\frac{3}{2}}|T|^{\frac{1}{2}}+|S|^{-\frac{1}{2}}S^*T|T|^{\frac{1}{2}} \\ |T|^{\frac{3}{2}}|S|^{\frac{1}{2}}+|T|^{-\frac{1}{2}}T^*S|S|^{\frac{1}{2}} & |T|^2+|T|^{-\frac{1}{2}}T^*S|T|^{\frac{1}{2}} \end{bmatrix}\right) \\ &\leq \| \, |S|+|T| \, \|.\end{aligned}$$

Thus, if S, T are positive operators, then

$$\begin{aligned}\|S+T\| &= \omega^{\frac{1}{2}}\left(\begin{bmatrix} S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}} & S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}} \\ T^{\frac{3}{2}}S^{\frac{1}{2}}+T^{\frac{1}{2}}S^{\frac{3}{2}} & T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}} \end{bmatrix}\right) \\ &= \left\|\begin{bmatrix} S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}} & S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}} \\ T^{\frac{3}{2}}S^{\frac{1}{2}}+T^{\frac{1}{2}}S^{\frac{3}{2}} & T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}} \end{bmatrix}\right\|^{\frac{1}{2}}.\end{aligned} \tag{2.5}$$

For positive invertible operators, (2.5) may be utilized to obtain the following upper bound for the norm of the sum of two such operators.

Corollary 2.7. *Let $S, T \in \mathbb{B}(\mathbb{H})$ be two positive operators. Then*

$$\begin{aligned}\|S+T\|^2 &\leq \frac{1}{2}\left(\left\|S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}}\right\|+\left\|T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}}\right\|\right) \\ &\quad + \frac{1}{2}\sqrt{\left(\left\|S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}}\right\|-\left\|T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}}\right\|\right)^2+4\left\|S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}}\right\|^2}.\end{aligned}$$

Proof. It follows from (2.5) and Lemma 1.4 that

$$\begin{aligned}&\|S+T\|^2 \\ &= \left\|\begin{bmatrix} S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}} & S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}} \\ T^{\frac{3}{2}}S^{\frac{1}{2}}+T^{\frac{1}{2}}S^{\frac{3}{2}} & T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}} \end{bmatrix}\right\| \\ &\leq \left\|\begin{bmatrix} \left\|S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}}\right\| & \left\|S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}}\right\| \\ \left\|T^{\frac{3}{2}}S^{\frac{1}{2}}+T^{\frac{1}{2}}S^{\frac{3}{2}}\right\| & \left\|T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}}\right\| \end{bmatrix}\right\| \\ &= \frac{1}{2}\left(\left\|S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}}\right\|+\left\|T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}}\right\|\right) \\ &\quad + \frac{1}{2}\sqrt{\left(\left\|S^2+S^{\frac{1}{2}}TS^{\frac{1}{2}}\right\|-\left\|T^2+T^{\frac{1}{2}}ST^{\frac{1}{2}}\right\|\right)^2+4\left\|S^{\frac{3}{2}}T^{\frac{1}{2}}+S^{\frac{1}{2}}T^{\frac{3}{2}}\right\|^2},\end{aligned}$$

where the last equality is obtained by direct calculations of the norm as the largest singular value of the matrix in context. This completes the proof. □

2.2. Upper bounds for the numerical radius

Related to the above discussion of the operator norm, we discuss some bounds for the numerical radius in the same context. In this section, it will be implicitly understood that the denominators are not zero by assumption.

Theorem 2.8. *Let $A_1, A_2, B_1, B_2 \in \mathbb{B}(\mathbb{H})$. Then*

$$\begin{aligned}&\omega(B_1A_1+B_2A_2) \\ &\leq \sqrt{\frac{\|A_1^*A_1+A_2^*A_2\| \, \|B_1B_1^*+B_2B_2^*\|}{\|A_1^*A_1+A_2^*A_2\|+\|B_1B_1^*+B_2B_2^*\|}}\|A_1^*A_1+A_2^*A_2+B_1B_1^*+B_2B_2^*\|^{\frac{1}{2}}.\end{aligned}$$

Proof. Let $0 < \nu < 1$. For any unit vector $x \in \mathbb{H}$, we have

$$\begin{aligned}
|\langle \mathbf{B}^*\mathbf{A}x, x\rangle| &= |\langle \mathbf{A}x, \mathbf{B}x\rangle| \\
&\leq \|\mathbf{A}x\|\,\|\mathbf{B}x\| \quad \text{(by the Cauchy-Schwarz inequality)} \\
&= \sqrt{\left\langle |\mathbf{A}|^2 x, x\right\rangle \left\langle |\mathbf{B}|^2 x, x\right\rangle} \\
&= \sqrt{\left\langle |\mathbf{A}|^{2\frac{\nu}{\nu}} x, x\right\rangle \left\langle |\mathbf{B}|^{2\frac{1-\nu}{1-\nu}} x, x\right\rangle} \\
&\leq \sqrt{\left\langle |\mathbf{A}|^{\frac{2}{\nu}} x, x\right\rangle^{\nu} \left\langle |\mathbf{B}|^{\frac{2}{1-\nu}} x, x\right\rangle^{1-\nu}} \quad \text{(by the Hölder-McCarthy inequality)} \\
&\leq \sqrt{\nu \left\langle |\mathbf{A}|^{\frac{2}{\nu}} x, x\right\rangle + (1-\nu)\left\langle |\mathbf{B}|^{\frac{2}{1-\nu}} x, x\right\rangle} \quad \text{(by the Young inequality)} \\
&= \sqrt{\left\langle \left(\nu|\mathbf{A}|^{\frac{2}{\nu}} + (1-\nu)\,|\mathbf{B}|^{\frac{2}{1-\nu}}\right) x, x\right\rangle} \\
&\leq \left\| \nu|\mathbf{A}|^{\frac{2}{\nu}} + (1-\nu)\,|\mathbf{B}|^{\frac{2}{1-\nu}} \right\|^{\frac{1}{2}}.
\end{aligned}$$

That is,

$$|\langle \mathbf{B}^*\mathbf{A}x, x\rangle| \leq \left\| \nu|\mathbf{A}|^{\frac{2}{\nu}} + (1-\nu)\,|\mathbf{B}|^{\frac{2}{1-\nu}} \right\|^{\frac{1}{2}}.$$

By taking the supremum over $x \in \mathbb{H}$ with $\|x\| = 1$, we get

$$\omega\left(\mathbf{B}^*\mathbf{A}\right) \leq \left\| \nu|\mathbf{A}|^{\frac{2}{\nu}} + (1-\nu)\,|\mathbf{B}|^{\frac{2}{1-\nu}} \right\|^{\frac{1}{2}}. \tag{2.6}$$

If we replace $\mathbf{A}$ and $\mathbf{B}$ by $\frac{\mathbf{A}}{\|\mathbf{A}\|}$ and $\frac{\mathbf{B}}{\|\mathbf{B}\|}$, in (2.6), we infer that

$$\omega\left(\frac{\mathbf{B}^*\mathbf{A}}{\|\mathbf{A}\|\,\|\mathbf{B}\|}\right) \leq \left\| \nu\left(\frac{|\mathbf{A}|^2}{\|\mathbf{A}\|^2}\right)^{\frac{1}{\nu}} + (1-\nu)\left(\frac{|\mathbf{B}|^2}{\|\mathbf{B}\|^2}\right)^{\frac{1}{1-\nu}} \right\|^{\frac{1}{2}}. \tag{2.7}$$

If we use the fact that if $O \leq X \leq I$, then $X^p \leq X$ for $p \geq 1$ (see [17]), we obtain from (2.7)

$$\omega\left(\frac{\mathbf{B}^*\mathbf{A}}{\|\mathbf{A}\|\,\|\mathbf{B}\|}\right) \leq \left\| \nu\frac{|\mathbf{A}|^2}{\|\mathbf{A}\|^2} + (1-\nu)\,\frac{|\mathbf{B}|^2}{\|\mathbf{B}\|^2} \right\|^{\frac{1}{2}}. \tag{2.8}$$

Now putting $\nu = \frac{\|\mathbf{A}\|^2}{\|\mathbf{A}\|^2+\|\mathbf{B}\|^2}$ and $1-\nu = \frac{\|\mathbf{B}\|^2}{\|\mathbf{A}\|^2+\|\mathbf{B}\|^2}$, in (2.8), we deduce that

$$\omega\left(\frac{\mathbf{B}^*\mathbf{A}}{\|\mathbf{A}\|\,\|\mathbf{B}\|}\right) \leq \left\| \frac{\|\mathbf{A}\|^2}{\|\mathbf{A}\|^2+\|\mathbf{B}\|^2}\frac{|\mathbf{A}|^2}{\|\mathbf{A}\|^2} + \frac{\|\mathbf{B}\|^2}{\|\mathbf{A}\|^2+\|\mathbf{B}\|^2}\frac{|\mathbf{B}|^2}{\|\mathbf{B}\|^2} \right\|^{\frac{1}{2}},$$

which is equivalent to

$$\omega\left(\mathbf{B}^*\mathbf{A}\right) \leq \frac{\|\mathbf{A}\|\,\|\mathbf{B}\|}{\sqrt{\|\mathbf{A}\|^2+\|\mathbf{B}\|^2}} \left\| |\mathbf{A}|^2 + |\mathbf{B}|^2 \right\|^{\frac{1}{2}}. \tag{2.9}$$

Let $\mathbf{A} = \begin{bmatrix} A_1 & O \\ A_2 & O \end{bmatrix}$ and $\mathbf{B} = \begin{bmatrix} B_1^* & O \\ B_2^* & O \end{bmatrix}$. Simple calculations reveal that

$$\omega\left(\mathbf{B}^*\mathbf{A}\right) = \omega\left(\begin{bmatrix} B_1 & B_2 \\ O & O \end{bmatrix}\begin{bmatrix} A_1 & O \\ A_2 & O \end{bmatrix}\right) = \omega\left(\begin{bmatrix} B_1A_1 + B_2A_2 & O \\ O & O \end{bmatrix}\right) = \omega\left(B_1A_1 + B_2A_2\right),$$

$$\begin{aligned}
\left\||\mathbf{A}|^2+|\mathbf{B}|^2\right\| &= \left\|\begin{bmatrix} A_1^* & A_2^* \\ O & O \end{bmatrix}\begin{bmatrix} A_1 & O \\ A_2 & O \end{bmatrix}+\begin{bmatrix} B_1 & B_2 \\ O & O \end{bmatrix}\begin{bmatrix} B_1^* & O \\ B_2^* & O \end{bmatrix}\right\| \\
&= \left\|\begin{bmatrix} A_1^*A_1+A_2^*A_2 & O \\ O & O \end{bmatrix}+\begin{bmatrix} B_1B_1^*+B_2B_2^* & O \\ O & O \end{bmatrix}\right\| \\
&= \left\|\begin{bmatrix} A_1^*A_1+A_2^*A_2+B_1B_1^*+B_2B_2^* & O \\ O & O \end{bmatrix}\right\| \\
&= \|A_1^*A_1+A_2^*A_2+B_1B_1^*+B_2B_2^*\|,
\end{aligned}$$

$$\begin{aligned}
\|\mathbf{A}\| &= \left\|\begin{bmatrix} A_1 & O \\ A_2 & O \end{bmatrix}\right\| \\
&= \left\|\begin{bmatrix} A_1^* & A_2^* \\ O & O \end{bmatrix}\begin{bmatrix} A_1 & O \\ A_2 & O \end{bmatrix}\right\|^{\frac{1}{2}} \\
&= \left\|\begin{bmatrix} A_1^*A_1+A_2^*A_2 & O \\ O & O \end{bmatrix}\right\|^{\frac{1}{2}} \\
&= \|A_1^*A_1+A_2^*A_2\|^{\frac{1}{2}},
\end{aligned}$$

and

$$\begin{aligned}
\|\mathbf{B}\| &= \left\|\begin{bmatrix} B_1^* & O \\ B_2^* & O \end{bmatrix}\right\| \\
&= \left\|\begin{bmatrix} B_1 & B_2 \\ O & O \end{bmatrix}\begin{bmatrix} B_1^* & O \\ B_2^* & O \end{bmatrix}\right\|^{\frac{1}{2}} \\
&= \left\|\begin{bmatrix} B_1B_1^*+B_2B_2^* & O \\ O & O \end{bmatrix}\right\|^{\frac{1}{2}} \\
&= \|B_1B_1^*+B_2B_2^*\|^{\frac{1}{2}}.
\end{aligned}$$

Now, the result follows directly from (2.9). □

The following corollary is an immediate consequence of Theorem 2.8, which improves the sub-multiplicative property of the operator norm.

Corollary 2.9. *Let $A, B \in \mathbb{B}(\mathbb{H})$ be nonzero operators. If B^*A is normal, then*

$$\|B^*A\| \leq \frac{\left\||A|^2+|B|^2\right\|^{\frac{1}{2}}}{\sqrt{\|A\|^2+\|B\|^2}}\ \|A\|\,\|B\|.$$

Remark 2.10. By the triangle inequality for the operator norm, we have

$$\left\||A|^2+|B|^2\right\|^{\frac{1}{2}} \leq \sqrt{\left\||A|^2\right\|+\left\||B|^2\right\|} = \sqrt{\|\,|A|\,\|^2+\|\,|B|\,\|^2} = \sqrt{\|A\|^2+\|B\|^2}, \quad (2.10)$$

i.e., $\frac{\left\||A|^2+|B|^2\right\|^{\frac{1}{2}}}{\sqrt{\|A\|^2+\|B\|^2}} \leq 1$. Therefore, Corollary 2.9 provides an improvement of the well-known inequality $\|B^*A\| \leq \|A\|\,\|B\|$.

Remark 2.11. Inequality (2.9) can also be written in the following arrangement, provided that $B^*A \neq O$,

$$\sqrt{\|A\|^2+\|B\|^2} \leq \frac{\|A\|\,\|B\|}{\omega(B^*A)}\left\||A|^2+|B|^2\right\|^{\frac{1}{2}},$$

which delivers a reverse for the inequality (2.10). Of course, if A and B are positive operators such that $B^{\frac{1}{2}}A^{\frac{1}{2}} \neq O$, then

$$\|A\|+\|B\| \leq \frac{\|A\|\,\|B\|}{\omega^2\left(B^{\frac{1}{2}}A^{\frac{1}{2}}\right)}\ \|A+B\|,$$

which presents a reverse for the triangle inequality.

Corollary 2.12. *Let $A_1, A_2, B_1, B_2 \in \mathbb{B}(\mathbb{H})$. Then*

$$\begin{aligned}&\omega(B_1A_1 + B_2A_2)\\ &\leq \sqrt{\frac{\|A_1^*A_1 + A_2^*A_2 + B_1B_1^* + B_2B_2^*\|}{\|A_1^*A_1 + A_2^*A_2\| + \|B_1B_1^* + B_2B_2^*\|}}\sqrt{\|A_1^*A_1 + A_2^*A_2\|\,\|B_1B_1^* + B_2B_2^*\|}.\end{aligned}$$

Proof. It follows from Theorem 2.8 that

$$\begin{aligned}&\|A_1^*A_1 + A_2^*A_2 + B_1B_1^* + B_2B_2^*\|\\ &\leq \|A_1^*A_1 + A_2^*A_2\| + \|B_1B_1^* + B_2B_2^*\|\\ &\quad \text{(by the triangle inequality for the operator norm)}\\ &\leq \frac{\|A_1^*A_1 + A_2^*A_2\|\,\|B_1B_1^* + B_2B_2^*\|}{\omega^2(B_1A_1 + B_2A_2)}\,\|A_1^*A_1 + A_2^*A_2 + B_1B_1^* + B_2B_2^*\|.\end{aligned}$$

The result follows directly from the above inequality. □

Remark 2.13. If we replace A_1, B_1, A_2, B_2 by $\sqrt{\frac{\|B_1\|}{\|A_1\|}}A_1$, $\sqrt{\frac{\|A_1\|}{\|B_1\|}}B_1$, $\sqrt{\frac{\|B_2\|}{\|A_2\|}}A_2$, and $\sqrt{\frac{\|A_2\|}{\|B_2\|}}B_2$, in Corollary 2.12, we obtain

$$\begin{aligned}&\omega(B_1A_1 + B_2A_2)\\ &\leq \sqrt{\frac{\left\|\frac{\|B_1\|}{\|A_1\|}A_1^*A_1 + \frac{\|B_2\|}{\|A_2\|}A_2^*A_2 + \frac{\|A_1\|}{\|B_1\|}B_1B_1^* + \frac{\|A_2\|}{\|B_2\|}B_2B_2^*\right\|}{\left\|\frac{\|B_1\|}{\|A_1\|}A_1^*A_1 + \frac{\|B_2\|}{\|A_2\|}A_2^*A_2\right\| + \left\|\frac{\|A_1\|}{\|B_1\|}B_1B_1^* + \frac{\|A_2\|}{\|B_2\|}B_2B_2^*\right\|}}\\ &\quad\times\sqrt{\left\|\frac{\|B_1\|}{\|A_1\|}A_1^*A_1 + \frac{\|B_2\|}{\|A_2\|}A_2^*A_2\right\|\left\|\frac{\|A_1\|}{\|B_1\|}B_1B_1^* + \frac{\|A_2\|}{\|B_2\|}B_2B_2^*\right\|}\\ &\leq \sqrt{\frac{\left\|\frac{\|B_1\|}{\|A_1\|}A_1^*A_1 + \frac{\|B_2\|}{\|A_2\|}A_2^*A_2 + \frac{\|A_1\|}{\|B_1\|}B_1B_1^* + \frac{\|A_2\|}{\|B_2\|}B_2B_2^*\right\|}{\left\|\frac{\|B_1\|}{\|A_1\|}A_1^*A_1 + \frac{\|B_2\|}{\|A_2\|}A_2^*A_2\right\| + \left\|\frac{\|A_1\|}{\|B_1\|}B_1B_1^* + \frac{\|A_2\|}{\|B_2\|}B_2B_2^*\right\|}}\\ &\quad\times\sqrt{\max\{\|A_1\|\,\|B_1\|, \|A_2\|\,\|B_2\|\} + \sqrt{\frac{\|B_1\|\,\|B_2\|}{\|A_1\|\,\|A_2\|}}\,\|A_1A_2^*\|}\\ &\quad\times\sqrt{\max\{\|A_1\|\,\|B_1\|, \|A_2\|\,\|B_2\|\} + \sqrt{\frac{\|A_1\|\,\|A_2\|}{\|B_1\|\,\|B_2\|}}\,\|B_1^*B_2\|}.\end{aligned}$$

Corollary 2.14. *Let $S, T \in \mathbb{B}(\mathbb{H})$. Then for any $0 \leq t \leq 1$,*

$$\begin{aligned}&\omega(S+T)\\ &\leq \sqrt{\frac{\left\||S|^{2t} + |T|^{2t} + |S^*|^{2(1-t)} + |T^*|^{2(1-t)}\right\|}{\left\||S|^{2t} + |T|^{2t}\right\| + \left\||S^*|^{2(1-t)} + |T^*|^{2(1-t)}\right\|}}\sqrt{\left\||S|^{2t} + |T|^{2t}\right\|\left\||S^*|^{2(1-t)} + |T^*|^{2(1-t)}\right\|}.\end{aligned}$$

Proof. Let $S = V|S|$ and $T = U|T|$ be the polar decomposition of S and T. By letting $A_1 = |S|^t$, $B_1 = V|S|^{1-t}$, $A_2 = |T|^t$, $B_2 = U|T|^{1-t}$, in Corollary 2.12, we reach to the desired result. □

Corollary 2.15. *Let $T \in \mathbb{B}(\mathbb{H})$. Then for any $0 \leq t \leq 1$,*

$$\omega(T) \leq \frac{1}{2}\sqrt{\frac{\left\||T|^{2t}+|T^*|^{2t}+|T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}{\left\||T|^{2t}+|T^*|^{2t}\right\|+\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}}\sqrt{\left\||T|^{2t}+|T^*|^{2t}\right\|\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}.$$

Proof. Letting $S = T^*$ in Corollary 2.14, we get

$$2\|\Re T\| \leq \sqrt{\frac{\left\||T|^{2t}+|T^*|^{2t}+|T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}{\left\||T|^{2t}+|T^*|^{2t}\right\|+\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}}\sqrt{\left\||T|^{2t}+|T^*|^{2t}\right\|\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}.$$

Substituting T by $e^{\mathrm{i}\theta}T$, in the above inequality, implies that

$$2\left\|\Re\left(e^{\mathrm{i}\theta}T\right)\right\| \leq \sqrt{\frac{\left\||T|^{2t}+|T^*|^{2t}+|T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}{\left\||T|^{2t}+|T^*|^{2t}\right\|+\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}}\sqrt{\left\||T|^{2t}+|T^*|^{2t}\right\|\left\||T|^{2(1-t)}+|T^*|^{2(1-t)}\right\|}.$$

The result follows by taking the supremum over $\theta \in \mathbb{R}$. □

Remark 2.16. The case $t = \frac{1}{2}$, namely

$$\omega(T) \leq \frac{1}{2}\left\||T|+|T^*|\right\|,$$

has been shown in [27, Ineq. (8)].

Author contributions. All the co-authors have contributed equally in all aspects of the preparation of this submission.

Conflict of interest statement. The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability. No data was used for the research described in the article.

References

[1] A. Abu-Omar, F. Kittaneh, *Numerical radius inequalities for $n \times n$ operator matrices*, Linear Algebra Appl. **468**, 18–26, 2015.

[2] A. Abu-Omar, F. Kittaneh, *Upper and lower bounds for the numerical radius with an application to involution operators*, Rocky Mountain J. Math. **45**(4), 1055–1065, 2015.

[3] D. Afraz, R. Lashkaripour, and M. Bakherad, *Further norm and numerical radius inequalities for sum of Hilbert space operators*, Filomat **38**(9), 3235–3242, 2024.

[4] E. Alizadeh and A. Farokhina, *some refinements of numerical radius inequalities for Hilbert space operators*, Jordan J. Math. Stat. **15**(1), 55-63, 2022.

[5] B. Al-Naddaf, A. Burqan, and F. Kittaneh, *Generalized numerical radius inequalities involving positive semidefinite block matrices*, J. Math. Inequal. **17**(4), 1363–1370, 2023.

[6] M. W. Alomari, M. Sababheh, C. Conde, and H. R. Moradi, *Generalized Euclidean Operator Radius*, Georgian Math. J. **31**(3), 369–380, 2024.

[7] M. Alomari, S. Sahoo and M. Bakherad, *Further numerical radius inequalities*, J. Math. Inequal. **16**(1), 307–326, 2022.

[8] M. Bakherad and F. Kittaneh, *Numerical radius inequalities involving commutators of G_1 operators*, Complex Anal. Oper. Theory **13**(4), 1557–1567, 2019.
[9] M. Bakherad and K. Shebrawi, *Upper bounds for numerical radius inequalities involving off-diagonal operator matrices*, Ann. Funct. Anal. **9**(3), 297–309, 2018.
[10] A. Benmakhlouf, O. Hirzallah, and F. Kittaneh, *On the p-numerical radii of Hilbert space operators*, Linear Multilinear Algebra **69**(15), 2813–2829, 2021.
[11] R. Bhatia, F. Kittaneh, *On the singular values of a product of operators*, SIAM J. Matrix Anal. Appl. **11**, 272–277, 1990.
[12] P. Bhunia, S.S. Dragomir, M.S. Moslehian, K. Paul, *Lectures on numerical radius inequalities*, Infosys Science Foundation Series in Mathematical Sciences, Springer Cham, (2022), XII+209 pp. https://doi.org/10.1007/978-3-031-13670-2
[13] P. Bhunia and K. Paul, *A-numerical radius: new inequalities and characterization of equalities*, Hacet. J. Math. Stat. **52**(5), 1254–1262, 2023.
[14] C. Conde, F. Kittaneh, H. R. Moradi, and M. Sababheh, *Numerical radii of operator matrices in terms of certain complex combinations of operators*, Georgian Math, J. **31**(4), 575–586, 2024.
[15] A. Frakis and F. Kittaneh, On *p*-numerical radius inequalities for commutators of operators, Quaest. Math. 2024. https://doi.org/10.2989/16073606.2024.2397562
[16] T. Furuta, *Invitation to Linear Operators: From Matrices to Bounded Linear Operators on a Hilbert Space*, CRC Press, 2001.
[17] T. Furuta, *Extension of the Furuta inequality and Ando-Hiai log majorization*, Linear Algebra Appl. **219**, 139–155, 1995.
[18] I. Gümüş, H. R. Moradi and M. Sababheh, *Operator inequalities via accretive transforms*, Hacettepe Journal of Mathematics and Statistics **53**(1), 40–52, 2024.
[19] K. E. Gustafson, D. K. M. Rao, *Numerical Range: The Field of Values of Linear Operators and Matrices*, Springer, New York, 1997.
[20] M. Hajmohamadi, R. Lashkaripour, and M. Bakherad, *Some generalizations of numerical radius on off-diagonal part of 2×2 operator matrices*, J. Math. Inequal. **12**(2), 447–457, 2018.
[21] O. Hirzallah, F. Kittaneh, and K. Shebrawi, *Numerical radius inequalities for commutators of Hilbert space operators*, Numer. Funct. Anal. Optim. **32**(7), 739–749, 2011.
[22] M. Hosseini, B. Moosavi, and H. R. Moradi, *An alternative estimate for the numerical radius of Hilbert space operators*, Math. Slovaca **70**(1), 233–237, 2020.
[23] M. Hosseini, M. E. Omidvar, *Some reverse and numerical radius inequalities*, Math. Slovaca **68**(5), 1121–1128, 2018.
[24] J. C. Hou, H.K. Du, *Norm inequalities of positive operator matrices*, Integral Equ. Oper. Theory **22**, 281–294, 1995.
[25] F. Kittaneh, *Norm inequalities for certain operator sums*, J. Funct. Anal. **143**, 337–348, 1997.
[26] F. Kittaneh, *Norm inequalities for sums of positive operators*, J. Oper. Theory **48**, 95–103, 2002.
[27] F. Kittaneh, *A numerical radius inequality and an estimate for the numerical radius of the Frobenius companion matrix*, Studia Math. **158**(1), 11–17, 2003.
[28] F. Kittaneh, *Numerical radius inequalities for Hilbert space operators*, Studia Math. **168**(1), 73–80, 2005.
[29] F. Kittaneh, H. R. Moradi, and M. Sababheh, *Sharper bounds for the numerical radius*, Linear Multilinear Algebra https://doi.org/10.1080/03081087.2023.2177248
[30] H. R. Moradi, M. Sababheh, *More accurate numerical radius inequalities II*, Linear Multilinear Algebra **69**(5), 921–933, 2021.

[31] M. E. Omidvar, H. R. Moradi, and K. Shebrawi, *Sharpening some classical numerical radius inequalities*, Oper. Matrices **12**(2), 407–416, 2018.
[32] H. Qiao, G. Hai, and E. Bai, *Some refinements of numerical radius inequalities for* 2×2 *operator matrices*, J. Math. Inequal. **16**(2), 425–444, 2022.
[33] M. Sababheh, *Numerical radius inequalities via convexity*, Linear Algebra Appl. **549**, 67–78, 2018.
[34] M. Sababheh, H. R. Moradi, *More accurate numerical radius inequalities (I)*, Linear Multilinear Algebra **69**(10), 1964–1973, 2021.
[35] M. Sababheh and H. R. Moradi, *Numerical radius of Kronecker product of matrices*, J. Appl. Anal. Comput. **13**(5), 2943–2954, 2023.
[36] A. Sheikhhosseini, M. Khosravi, and M. Sababheh, *The weighted numerical radius*, Ann. Funct. Anal. **13**, Article 3, 2022.
[37] S. Sheybani, M. Sababheh, and H. R. Moradi, *Weighted inequalities for the numerical radius*, Vietnam J. Math. **51**(2), 363–377, 2023.
[38] S. Soltani, A. Frakis, *Further refinements of some numerical radius inequalities for operators*, Oper. Matrices **17**(1), 245–257, 2023.